



A modified Halpern-type iteration algorithm for a family of hemi-relatively nonexpansive mappings and systems of equilibrium problems in Banach spaces[☆]

Ziming Wang^a, Yongfu Su^{a,*}, Dongxing Wang^b, Yucai Dong^b

^a Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, PR China

^b Institute of Nonlinear Science, Academy of Armored Forced Engineering Institute, Beijing 100072, PR China

ARTICLE INFO

Article history:

Received 20 June 2009

Received in revised form 4 December 2009

MSC:

47H05

47H09

47H10

Keywords:

Hemi-relatively nonexpansive mappings

Hybrid algorithm

Systems of equilibrium problems

Fixed point

Banach space

ABSTRACT

In this paper, we prove strong convergence theorems by the hybrid method for a family of hemi-relatively nonexpansive mappings in a Banach space. Our results improve and extend the corresponding results given by Qin et al. [Xiaolong Qin, Yeol Je Cho, Shin Min Kang, Haiyun Zhou, Convergence of a modified Halpern-type iteration algorithm for quasi- ϕ -nonexpansive mappings, *Appl. Math. Lett.* 22 (2009) 1051–1055], and at the same time, our iteration algorithm is different from the Kimura and Takahashi algorithm, which is a modified Mann-type iteration algorithm [Yasunori Kimura, Wataru Takahashi, On a hybrid method for a family of relatively nonexpansive mappings in Banach space, *J. Math. Anal. Appl.* 357 (2009) 356–363]. In addition, we succeed in applying our algorithm to systems of equilibrium problems which contain a family of equilibrium problems.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Let E be a real Banach space, and let E^* be the dual space of E . Let C be a nonempty closed convex subset of E . Let f be a bifunction of $C \times C$ into \mathbb{R} , where \mathbb{R} is the set of real numbers. The equilibrium problem for $f : C \times C \rightarrow \mathbb{R}$ is to find $x \in C$ such that

$$f(x, y) \geq 0 \quad \text{for all } y \in C. \quad (1.1)$$

The set of solutions of (1.1) is denoted by $EP(f)$. Given a mapping $T : C \rightarrow H$, let $f(x, y) = \langle Tx, y - x \rangle$ for all $x, y \in C$. Then, $z \in EP(f)$ if and only if $\langle Tz, y - z \rangle \geq 0$ for all $y \in C$, i.e., z is a solution of the variational inequality.

Equilibrium problems, which were introduced in [1] in 1994, have had a great impact and influence in the development of several branches of pure and applied sciences. It has been shown that equilibrium problem theory provides a novel and unified treatment of a wide class of problems which arise in economics, finance, physics, image reconstruction, ecology, transportation, network, elasticity, and optimization. Numerous problems in physics, optimization, and economics reduce to finding a solution of (1.1). Some methods have been proposed to solve the equilibrium problem; see, for instance, [1–14] and the references therein.

[☆] Supported by the National Natural Science Foundation of China under grant (10771050).

* Corresponding author.

E-mail addresses: ziming.wang@hotmail.com (Z. Wang), suyongfu@tjpu.edu.cn (Y. Su).

A mapping $T : C \rightarrow C$ is said to be nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for every $x, y \in C$. Iterative methods for the approximation of fixed points of a nonexpansive mapping have been studied by many researchers; see, for example, [4–6,8,15,16].

On the other hand, the hemi-relatively nonexpansive mapping, which is another generalization of a nonexpansive mapping and a relatively nonexpansive mapping, has been considered recently. Its properties and the iterative schemes for such a mapping have been studied in [12,17,18], amongst others.

Recently, Qin et al. [17] established the strong convergence of an iterative scheme with a new type of hybrid method as follows.

Theorem 1.1 (See Qin et al. [17]). *Let C be a nonempty and closed convex subset of a uniformly convex and uniformly smooth Banach space E , and let $T : C \rightarrow C$ be a closed and hemi-relatively nonexpansive mapping such that $F(T) \neq \emptyset$. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{cases} x_0 \in E & \text{chosen arbitrarily,} \\ C_1 = C, \\ x_1 = \Pi_{C_1} x_0, \\ y_n = J^{-1}(\alpha_n x_1 + (1 - \alpha_n)Tx_n), \\ C_{n+1} = \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x_1, \quad \forall n \geq 1. \end{cases} \quad (1.2)$$

Assume that the control sequence satisfies the restriction $\lim_{n \rightarrow \infty} \alpha_n = 0$. Then $\{x_n\}$ converges strongly to $\Pi_{F(T)} x_1$.

Motivated by this result, we prove strong convergence theorems by the hybrid method for a family of hemi-relatively nonexpansive mappings in a Banach space. The main result is more general than Theorem 1.1, and at the same time, our iteration algorithm is different from the Kimura and Takahashi algorithm, which is a modified Mann-type iteration algorithm, given in [19]. In addition, we succeed in applying our algorithm to systems of equilibrium problems which contain a family of equilibrium problems.

2. Preliminaries

In what follows, E denotes a real Banach space with norm $\|\cdot\|$ and E^* the dual space of E . The norm of E^* is also denoted by $\|\cdot\|$. For $y^* \in E^*$, its value at $x \in E$ is denoted by $\langle x, y^* \rangle$.

A Banach space E is said to be strictly convex if, for $x, y \in E$, $\|x\| = \|y\| = 1$, $x \neq y$ implies that $\|x + y\| < 2$. E is said to have the Kadec–Klee property if a weakly convergent sequence $\{x_n\}$ in E with limit $x_0 \in E$ satisfies that $\lim_{n \rightarrow \infty} \|x_n\| = \|x_0\|$; then $\{x_n\}$ converges strongly to x_0 . Let $S_E = \{x \in E : \|x\| = 1\}$, and define $f : S_E \times S_E \times \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ by

$$f(x, y, t) = \frac{\|x + ty\| - \|x\|}{t}$$

for $x, y \in S_E$ and $t \in \mathbb{R} \setminus \{0\}$. A norm of E is said to be Gâteaux differentiable if $\lim_{t \rightarrow 0} f(x, y, t)$ has a limit for each $x, y \in S_E$. In this case, E is said to be smooth. A norm of E is said to be Fréchet differentiable if $\lim_{t \rightarrow 0} f(x, y, t)$ is attained uniformly for $y \in S_E$ for each $x \in E$. It is known that E^* has a Fréchet differentiable norm if and only if E is strictly convex and reflexive, and has the Kadec–Klee property.

The normalized duality mapping of E is denoted by J ; that is,

$$Jx = \{x^* \in E^* : \|x\|^2 = \langle x, x^* \rangle = \|x^*\|^2\}$$

for $x \in E$. If E is a smooth, strictly convex, and reflexive Banach space, then J is a single-valued one-to-one mapping onto E^* . In this case, the inverse mapping J^{-1} coincides with the duality mapping on E . For more details, see [20].

Suppose that a Banach space E is smooth. Then J is a single-valued mapping. The function $\phi : E \times E \rightarrow \mathbb{R}$ is defined by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2$$

for $x, y \in E$. We know several fundamental properties of ϕ , as follows. $\phi(x, y) \geq 0$ for all $x, y \in E$. For a sequence $\{y_n\}$ in E and $x \in E$, $\{y_n\}$ is bounded if and only if $\{\phi(x, y_n)\}$ is bounded. For more details, see, for example, [5].

Let C be a closed convex subset of E , and let T be a mapping from C into itself. A point p in C is said to be an asymptotic fixed point of T if C contains a sequence $\{x_n\}$ which converges weakly to p such that the strong limit $\lim_{n \rightarrow \infty} (x_n - Tx_n) = 0$. The set of asymptotic fixed points of T will be denoted by $\widehat{F}(T)$. A mapping T from C into itself is called a nonexpansive mapping if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$ and a relatively nonexpansive mapping [21–23] if $\widehat{F}(T) = F(T)$ and $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$. The asymptotic behavior of a relatively nonexpansive mapping was studied in [21–23]. A point p in C is said to be a strong asymptotic fixed point of T if C contains a sequence $\{x_n\}$ which converges strongly to p such that $\lim_{n \rightarrow \infty} (x_n - Tx_n) = 0$. The set of strong asymptotic fixed points of T will be denoted by $\widetilde{F}(T)$. A mapping T from C into itself is called a relatively weak nonexpansive mapping if $\widetilde{F}(T) = F(T)$ and $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$. A mapping T is called a hemi-relatively nonexpansive mapping if $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$.

It is obvious that a relatively nonexpansive mapping is a relatively weak nonexpansive mapping, and a relatively weak nonexpansive mapping is a hemi-relatively nonexpansive mapping; however, the converse is not true. In order to explain this better, we give the following example.

Example 2.1 ([18]). Let E be any smooth Banach space, and let $x_0 \neq 0$ be any element of E . We define a mapping $T : E \rightarrow E$ as follows:

$$T(x) = \begin{cases} \left(\frac{1}{2} + \frac{1}{2^n}\right)x_0, & \text{if } x = \left(\frac{1}{2} + \frac{1}{2^n}\right)x_0, \\ -x, & \text{if } x \neq \left(\frac{1}{2} + \frac{1}{2^n}\right)x_0, \end{cases}$$

for $n = 1; 2; 3; \dots$. Then T is a hemi-relatively nonexpansive mapping but not a relatively nonexpansive mapping.

Next, we give some important examples which are hemi-relatively nonexpansive.

Example 2.2 ([10]). Let E be a strictly convex reflexive smooth Banach space. Let A be a maximal monotone operator of E into E^* and let J_r be the resolvent for A with $r > 0$. Then $J_r = (J + rA)^{-1}J$ is a hemi-relatively nonexpansive mapping from E onto $D(A)$ with $F(J_r) = A^{-1}0$.

Remark 2.3. There are other examples of hemi-relatively nonexpansive mappings such as the generalized projections (or projections), and others; see [10].

Let $\{C_n\}$ be a sequence of nonempty closed convex subsets of a reflexive Banach space E . We define two subsets $s - Li_n C_n$ and $w - Ls_n C_n$ as follows: $x \in s - Li_n C_n$ if and only if there exists $\{x_n\} \subset E$ such that $\{x_n\}$ converges strongly to x and such that $x_n \in C_n$ for all $n \in \mathbb{N}$. Similarly, $y \in w - Ls_n C_n$ if and only if there exists a subsequence $\{C_{n_i}\}$ of $\{C_n\}$ and a sequence $\{y_i\} \subset E$ such that $\{y_i\}$ converges weakly to y and such that $y_i \in C_{n_i}$ for all $i \in \mathbb{N}$. We define the Mosco convergence [24] of $\{C_n\}$ as follows. If C_0 satisfies that $C_0 = s - Li_n C_n = w - Ls_n C_n$, it is said that $\{C_n\}$ converges to C_0 in the sense of Mosco, and we write $C_0 = M - \lim_{n \rightarrow \infty} C_n$. For more details, see [25].

Let C be a nonempty closed convex subset of a smooth, strictly convex, and reflexive Banach space E . Then, for arbitrarily fixed $x \in E$, a function $C \ni \|y - x\| \in \mathbb{R}$ has a unique minimizer $y_x \in C$. Using such a point, we define the metric projection P_C by $P_C x = y_x = \arg \min_{y \in C} \|x - y\|$ for every $x \in E$. In a similar fashion, we can see that a function $C \ni y \mapsto \phi(x, y) \in \mathbb{R}$ also has a unique minimizer $z_x \in C$. The generalized projection Π_C of E onto C is defined by $\Pi_C x = z_x = \arg \min_{y \in C} \phi(x, y)$ for every $x \in E$; see [1].

In order to prove our results, the following theorem plays an important role.

Theorem 2.4 ([26]). Let E be a smooth, reflexive, and strictly convex Banach space having the Kadec–Klee property. Let $\{K_n\}$ be a sequence of nonempty closed convex subsets of E . If $K_0 = M - \lim_{n \rightarrow \infty} K_n$ exists and is nonempty, then $\{\Pi_{K_n} x\}$ converges strongly to $\Pi_{K_0} x$ for each $x \in C$.

3. A modified Halpern-type iteration algorithm for a family of hemi-relatively nonexpansive mappings

First, let us consider a family of hemi-relatively nonexpansive mappings $\{S_\lambda : \lambda \in \Lambda\}$ defined on a nonempty closed convex subset of a Banach space, and suppose that there exists a common fixed point. We prove that an iterative scheme generated by $\{S_\lambda\}$ converges strongly to a point of the set $\bigcap_\lambda F(S_\lambda)$ of common fixed points of $\{S_\lambda\}$. The iterative scheme we adopt is originally defined in [18].

The following lemma is also important in our main results.

Lemma 3.1. Let E be a strictly convex reflexive Banach space having a Fréchet differentiable norm, C a nonempty closed convex subset of E , and $\{S_i\}$ a sequence of mappings of C into itself. Let $\{x_i\}$ be a strongly convergent sequence in C with a limit x_0 and $\{y_i\}$ a sequence in C defined by $y_i = J^{-1}(\alpha_i x_1 + (1 - \alpha_i) S_i x_i)$ for each $i \in \mathbb{N}$, where $\{\alpha_i\}$ is a convergent sequence in $[0, 1]$ with $\lim_{i \rightarrow \infty} \alpha_i = 0$. Suppose that $\phi(x_0, y_i) \leq \phi(x_0, x_i)$ for all $i \in \mathbb{N}$ and that $\{y_i\}$ converges weakly to $y_0^* \in E^*$. Then, $\{x_i - J S_i x_i\}$ converges strongly to 0. Moreover, if E has the Kadec–Klee property, then $\{S_i x_i\}$ converges strongly to x_0 .

Proof. Since $\phi(x_0, y_i) \leq \phi(x_0, x_i)$ for all $i \in \mathbb{N}$, we have that

$$0 \leq \lim_{i \rightarrow \infty} \phi(x_0, y_i) \leq \lim_{i \rightarrow \infty} \phi(x_0, x_i) = 0,$$

and hence $\lim_{i \rightarrow \infty} \phi(x_0, y_i) = 0$. Since

$$(\|x_0\| - \|y_i\|)^2 = \|x_0\|^2 - 2\|x_0\| \|y_i\| + \|y_i\|^2 \leq \phi(x_0, y_i)$$

for $i \in \mathbb{N}$, we obtain that $\lim_{i \rightarrow \infty} \|y_i\| = \|x_0\|$ and that

$$\lim_{i \rightarrow \infty} \langle x_0, y_i \rangle = \lim_{i \rightarrow \infty} \frac{1}{2} (\|x_0\|^2 + \|y_i\|^2 - \phi(x_0, y_i)) = \|x_0\|^2.$$

Using weak lower semicontinuity of the norm, we have that

$$\begin{aligned}\|x_0\|^2 &= \lim_{i \rightarrow \infty} \langle x_0, Jy_i \rangle = \langle x_0, y_0^* \rangle \leq \|x_0\| \|y_0^*\| \\ &\leq \|x_0\| \liminf_{i \rightarrow \infty} \|Jy_i\| = \|x_0\| \lim_{i \rightarrow \infty} \|Jy_i\| \\ &= \|x_0\|^2.\end{aligned}$$

Therefore, we have that $\|y_0^*\|^2 = \langle x_0, y_0^* \rangle = \|x_0\|^2$ and hence $y_0^* = Jx_0$. Thus we have that $\{Jy_i\}$ converges weakly to Jx_0 . It also holds that

$$\lim_{i \rightarrow \infty} \|Jy_i\| = \lim_{i \rightarrow \infty} \|y_i\| = \|x_0\| = \|Jx_0\|.$$

Since E has a Fréchet differentiable norm, it follows that E^* has the Kadec–Klee property, and thus we have that $\{Jy_i\}$ converges strongly to Jx_0 . Then, we have that

$$\begin{aligned}\|Jx_0 - Jy_i\| &= \|Jx_0 - (\alpha_i Jx_1 + (1 - \alpha_i)JS_i x_i)\| \\ &\geq \|Jx_0 - \alpha_i Jx_0 - (1 - \alpha_i)JS_i x_i\| - \alpha_i \|Jx_1 - Jx_0\| \\ &= (1 - \alpha_i)\|Jx_0 - JS_i x_i\| - \alpha_i \|Jx_1 - Jx_0\|\end{aligned}$$

for $i \in \mathbb{N}$. Using norm-to-norm continuity of J , and $\alpha_i \rightarrow 0$, we get that

$$\lim_{i \rightarrow \infty} [(1 - \alpha_i)\|Jx_0 - JS_i x_i\| - \alpha_i \|Jx_1 - Jx_0\|] = \lim_{n \rightarrow \infty} \|Jx_0 - JS_i x_i\| = 0.$$

We also have that $\{x_i\}$ converges strongly to Jx_0 , and hence we obtain that $\{x_i - JS_i x_i\}$ converges strongly to 0. Further, let us suppose that E has the Kadec–Klee property. Then, the norm of E^* is Fréchet differentiable, and therefore J^{-1} is norm-to-norm continuous. Hence, we have that

$$\lim_{i \rightarrow \infty} \|x_0 - Sx_i\| = \lim_{i \rightarrow \infty} \|J^{-1}Jx_0 - J^{-1}JS_i x_i\| = 0,$$

which is the desired result. \square

Using this lemma, we obtain the following convergence theorem for a modified Halpern-type iteration algorithm generated by a family of hemi-relatively nonexpansive mappings.

Theorem 3.2. *Let E be a strictly convex reflexive Banach space having the Kadec–Klee property and a Fréchet differentiable norm. Let C be a nonempty closed convex subset of E and $\{S_\lambda : \lambda \in \Lambda\}$ a family of closed hemi-relatively nonexpansive mappings of C into itself having a common fixed point. Let $\{\alpha_n\}$ be a sequence in $[0, 1]$ such that $\lim_{n \rightarrow \infty} \alpha_n = 0$. For an arbitrarily chosen point $x \in E$, generate a sequence $\{x_n\}$ by the following iterative scheme:*

$$\begin{cases} x_1 \in C & \text{chosen arbitrarily,} \\ C_1 = C, \\ y_{n,\lambda} = J^{-1}(\alpha_n Jx_1 + (1 - \alpha_n)JS_\lambda x_n), \\ C_{n+1} = \{z \in C_n : \sup_{\lambda \in \Lambda} \phi(z, y_{n,\lambda}) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x \end{cases} \quad (3.1)$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to $\Pi_F x$, where Π_F is the generalized projection from C onto F , and $F = \bigcap_{\lambda \in \Lambda} F(S_\lambda)$ is the set of common fixed points of $\{S_\lambda\}$.

Proof. First, we show that C_n is closed and convex. In view of the definition of ϕ , we can obtain that

$$\begin{aligned}C_{n+1} &= \{z \in C_n : \sup_{\lambda \in \Lambda} \phi(z, y_{n,\lambda}) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\} \\ &= \bigcap_{\lambda \in \Lambda} \{z \in C_n : \phi(z, y_{n,\lambda}) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\} \\ &= \bigcap_{\lambda \in \Lambda} \{z \in C : 2\alpha_n \langle z, Jx_1 \rangle + 2(1 - \alpha_n)\langle z, Jx_n \rangle \\ &\quad - 2\langle z, Jy_{n,\lambda} \rangle - \alpha_n \|x_1\|^2 - (1 - \alpha_n)\|x_n\|^2 + \|y_{n,\lambda}\|^2 \leq 0\} \cap C_n,\end{aligned} \quad (3.2)$$

which yields that $\{C_n\}$ is closed and convex for every $n \in \mathbb{N}$.

Second, we prove that $F = \bigcap_{\lambda \in \Lambda} F(S_\lambda) \subset C_n$, for all $n \in \mathbb{N}$. Let $p \in \bigcap_{\lambda \in \Lambda} F(S_\lambda)$. Then, for each $n \in \mathbb{N}$ and $\lambda \in \Lambda$, we have that

$$\begin{aligned}\phi(p, y_{n,\lambda}) &= \phi(p, J^{-1}(\alpha_n Jx_1 + (1 - \alpha_n)JS_\lambda x_n)) \\ &= \|p\|^2 - 2\langle p, \alpha_n Jx_1 + (1 - \alpha_n)JS_\lambda x_n \rangle + \|\alpha_n Jx_1 + (1 - \alpha_n)JS_\lambda x_n\|^2\end{aligned}$$

$$\begin{aligned}
&\leq \|p\|^2 - 2\alpha_n \langle p, Jx_1 \rangle - 2(1 - \alpha_n) \langle p, JS_\lambda x_n \rangle + \alpha_n \|x_1\|^2 + (1 - \alpha_n) \|S_\lambda x_n\|^2 \\
&= \alpha_n (\|p\|^2 - 2\langle p, Jx_1 \rangle + \alpha_n \|x_1\|^2) + (1 - \alpha_n) (\|p\|^2 - 2\langle p, JS_\lambda x_n \rangle + \|S_\lambda x_n\|^2) \\
&\leq \alpha_n \phi(p, x_1) + (1 - \alpha_n) \phi(p, x_n).
\end{aligned} \tag{3.3}$$

Therefore, $p \in C_n$ for all $n \in \mathbb{N}$, and hence $F = \bigcap_{\lambda \in \Lambda} F(S_\lambda) \subset C_n$ for all $n \in \mathbb{N}$. Since F is nonempty, C_n is a nonempty closed convex subset of E , and thus Π_{C_n} exists for every $n \in \mathbb{N}$. Hence $\{x_n\}$ is well defined.

Third, we shall show that $\lim_{n \rightarrow \infty} x_n = x_0 = P_{C_0}x$. Since $\{C_n\}$ is a decreasing sequence of closed convex subsets of E such that $C_0 = \bigcap_{n=1}^{\infty} C_n$ is nonempty, it follows that

$$M - \lim_{n \rightarrow \infty} C_n = C_0 = \bigcap_{n=1}^{\infty} C_n \neq \emptyset.$$

By Theorem 2.4, $\{x_n\} = \{P_{C_n}x\}$ converges strongly to $x_0 = P_{C_0}x$.

Fourth, we prove that $x_0 \in F = \bigcap_{\lambda \in \Lambda} F(S_\lambda)$. Since $x_0 \in C_n$ for every $n \in \mathbb{N}$, it follows that

$$\sup_{\lambda \in \Lambda} \phi(x_0, y_{n,\lambda}) \leq \alpha_n \phi(x_0, x_1) + (1 - \alpha_n) \phi(x_0, x_n)$$

for all $n \in \mathbb{N}$. Fix $\lambda \in \Lambda$ arbitrarily. From the assumption that $\lim_{n \rightarrow \infty} \alpha_n = 0$, we can take a subsequence $\{y_{n_i,\lambda}\}$ of $\{y_{n,\lambda}\}$ such that $\{y_{n_i,\lambda}\}$ converges weakly to a point $y^* \in E^*$. Then, by Lemma 3.1, we have that

$$\lim_{i \rightarrow \infty} \|x_0 - S_\lambda x_{n_i}\| = 0.$$

And from the closedness of S_λ , for any $\lambda \in \Lambda$, we have that $x_0 \in F(S_\lambda)$ for any $\lambda \in \Lambda$, and hence $x_0 \in F = \bigcap_{\lambda \in \Lambda} F(S_\lambda)$.

Finally, since $x_0 = \Pi_{C_0}x \in F$ and F is a nonempty closed convex subset of $C_0 = \bigcap_{n=1}^{\infty} C_n$, we conclude that $x_0 = \Pi_F x$, which completes the proof. \square

Remark 3.3.

- (1) We extend the main result of [17] from a hemi-relatively nonexpansive mapping to a general family of hemi-relatively nonexpansive mappings; furthermore, our method of proof is different from that in [17].
- (2) Our iterative scheme is a modified Halpern-type scheme. In [19], the authors studied a modified Mann-type scheme; they are independent of each other.

Considering a single operator as a special case, we can obtain the following corollary directly.

Corollary 3.4. Let E be a strictly convex reflexive Banach space having the Kadec–Klee property and a Fréchet differentiable norm. Let C be a nonempty closed convex subset of E and S a closed hemi-relatively nonexpansive mappings of C into itself having a common fixed point. Let $\{\alpha_n\}$ be a sequence in $[0, 1]$ such that $\lim_{n \rightarrow \infty} \alpha_n = 0$. For an arbitrarily chosen point $x \in E$, generate a sequence $\{x_n\}$ by the following iterative scheme:

$$\begin{cases} x_1 \in C & \text{chosen arbitrarily,} \\ C_1 = C, \\ y_n = J^{-1}(\alpha_n Jx_1 + (1 - \alpha_n)JSx_n), \\ C_{n+1} = \{z \in C_n : \phi(z, y_n) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}}x \end{cases} \tag{3.4}$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to $\Pi_{F(S)}x$, where $\Pi_{F(S)}$ is the generalized projection from C onto $F(S)$.

4. A modified Halpern-type iteration algorithm for systems of equilibrium problems

In this section, we consider the problem of finding a solution of EP . Equilibrium problems provide a novel and unified treatment of a wide class of problems which arise in economics, finance, physics, image reconstruction, ecology, transportation, network, elasticity, optimization, and so on.

For solving the equilibrium problem for a bifunction $f : C \times C \rightarrow \mathbb{R}$, let us assume that f satisfies the following conditions:

- (A₁) $f(x, x) = 0$, for all $x \in C$;
- (A₂) f is monotone, i.e., $f(x, y) + f(y, x) \leq 0$, for all $x, y \in C$;
- (A₃) for each $x, y, z \in C$, $\lim_{t \downarrow 0} f(tz + (1 - t)x, y) \leq f(x, y)$;
- (A₄) for each $x \in C$, the function $y \mapsto f(x, y)$ is convex and lower semicontinuous.

Lemma 4.1 ([1]). Let C be a nonempty closed convex subset of a smooth, strictly convex, and reflexive Banach space E , and let $f : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A₁)–(A₄). Let $r > 0$ and $x \in E$; then, there exists $z \in C$ such that

$$f(z, y) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0, \quad \text{for all } y \in C.$$

Lemma 4.2 ([13]). Let C be a nonempty closed convex subset of a uniformly smooth, strictly convex, and reflexive Banach space E , and let $f : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A_1) – (A_4) . Let $r > 0$ and $x \in E$, and define a mapping $T_r : E \rightarrow C$ as follows:

$$T_r x = \left\{ z \in C : f(z, y) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0, \forall y \in C \right\}$$

for all $z \in C$. Then, the following hold:

1. T_r is single-valued;
2. T_r is firmly nonexpansive, i.e., for any $x, y \in H$,

$$\langle T_r x - T_r y, JT_r x - JT_r y \rangle \leq \langle T_r x - T_r y, Jx - Jy \rangle;$$
3. $F(T_r) = EP(f)$;
4. $EP(f)$ is closed and convex.

Lemma 4.3 ([14]). Let C be a closed convex subset of a smooth, strictly convex and reflexive Banach space E , let $f : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A_1) – (A_4) , and let $r > 0$. Then, for $x \in E$ and $q \in F(T_r)$,

$$\phi(q, T_r x) + \phi(T_r x, x) \leq \phi(q, x).$$

In order to use the iterative scheme in Theorem 3.2 to deal with equilibrium problems, we give the following conclusion, which has studied by some authors (see [10,27]).

Lemma 4.4. Let C be a closed convex subset of uniformly smooth, strictly convex and reflexive Banach space E , and define the mapping T_r as Lemma 4.2; then T_r is a hemi-relatively nonexpansive mapping.

Next, we consider the equilibrium problem for a family of bifunctions $\{f_\lambda : C \times C \rightarrow \mathbb{R}, \lambda \in \Lambda\}$, which satisfy (A_1) – (A_4) . Very recently, Colao, Aceto, and Marino have studied a similar case. The difference is that $\lambda \in \mathbb{N}$ in their case, and the case is said to be that of systems of equilibrium problems; see [28].

In the following, we will apply the iterative scheme (3.1) to systems of equilibrium problems.

Theorem 4.5. Let E be a strictly convex reflexive Banach space having the Kadec–Klee property and a Fréchet differentiable norm. Let C be a nonempty closed convex subset of E . Let $f_\lambda : C \times C \rightarrow \mathbb{R}, \lambda \in \Lambda$ be a family of bifunctions satisfying (A_1) – (A_4) , and $F = \bigcap_{\lambda \in \Lambda} EP(f_\lambda) \neq \emptyset$. Suppose that $\{\alpha_n\}$ is a sequence in $[0, 1]$ such that $\lim_{n \rightarrow \infty} \alpha_n = 0$. For an arbitrarily chosen point $x \in E$, generate a sequence $\{x_n\}$ by the following iterative scheme:

$$\begin{cases} x_1 \in C \text{ chosen arbitrarily,} \\ C_1 = C, \\ f_\lambda(u_n, y) + \frac{1}{r} \langle y - u_n, Ju_n - Jx_n \rangle \geq 0, \quad \forall y \in C, r > 0, \\ y_{n,\lambda} = J^{-1}(\alpha_n Jx_1 + (1 - \alpha_n)Ju_{n,\lambda}), \\ C_{n+1} = \{z \in C_n : \sup_{\lambda \in \Lambda} \phi(z, y_{n,\lambda}) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x \end{cases} \quad (4.1)$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to $\Pi_F x$, where $F = \bigcap_{\lambda \in \Lambda} EP(f_\lambda)$.

Proof. In view of Lemma 4.2, we know that $u_{n,\lambda} = T_{r,\lambda} x_n$ and $F(T_{r,\lambda}) = EP(f_\lambda)$ for all $\lambda \in \Lambda$. So we can replace the iterative scheme (4.1) by the following:

$$\begin{cases} x_1 \in C \text{ chosen arbitrarily,} \\ C_1 = C, \\ y_{n,\lambda} = J^{-1}(\alpha_n Jx_1 + (1 - \alpha_n)JT_{r,\lambda} x_n), \\ C_{n+1} = \{z \in C_n : \sup_{\lambda \in \Lambda} \phi(z, y_{n,\lambda}) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x. \end{cases} \quad (4.2)$$

From Lemma 4.4, we obtain that $T_{r,\lambda}$ must be a hemi-relatively nonexpansive mapping for $\lambda \in \Lambda$. And by Lemma 4.2 again, we learn that $T_{r,\lambda}$ is a family of continuous operators. So $T_{r,\lambda}$ must be closed. Using Theorem 3.2, we can get that $\{x_n\}$ converges strongly to $\Pi_{F_1} x$, where $F_1 = \bigcap_{\lambda \in \Lambda} T_{r,\lambda}$, and by Lemma 4.2, $F_1 = \bigcap_{\lambda \in \Lambda} EP(f_\lambda) = F$, which completes the proof. \square

As a special case, we consider a single $EP(f)$, and we can directly give the following corollary.

Corollary 4.6. Let E be a strictly convex reflexive Banach space having the Kadec–Klee property and a Fréchet differentiable norm. Let C be a nonempty closed convex subset of E . Let $f : C \times C \rightarrow \mathbb{R}$, $\lambda \in \Lambda$ be a bifunction satisfying (A_1) – (A_4) , and $EP(f) \neq \emptyset$. Suppose that $\{\alpha_n\}$ is a sequence in $[0, 1]$ such that $\lim_{n \rightarrow \infty} \alpha_n = 0$. For an arbitrarily chosen point $x \in E$, generate a sequence $\{x_n\}$ by the following iterative scheme:

$$\begin{cases} x_1 \in C \text{ chosen arbitrarily,} \\ C_1 = C, \\ f(u_n, y) + \frac{1}{r} \langle y - u_n, Ju_n - Jx_n \rangle \geq 0, \quad \forall y \in C, r > 0, \\ y_n = J^{-1}(\alpha_n Jx_1 + (1 - \alpha_n)Ju_n), \\ C_{n+1} = \{z \in C_n : \phi(z, y_n) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x \end{cases} \quad (4.3)$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to $\Pi_{EP(f)} x$.

If we consider another case, that is, $\alpha_n \equiv 0$ in (4.1), we can apply Theorem 4.5 and obtain the following corollary directly.

Corollary 4.7. Let E be a strictly convex reflexive Banach space having the Kadec–Klee property and a Fréchet differentiable norm. Let C be a nonempty closed convex subset of E . Let $f_\lambda : C \times C \rightarrow \mathbb{R}$, $\lambda \in \Lambda$ be a family of bifunctions satisfying (A_1) – (A_4) , and $F = \bigcap_{\lambda \in \Lambda} EP(f_\lambda) \neq \emptyset$. Suppose that $\alpha_n \equiv 0$. For an arbitrarily chosen point $x \in E$, generate a sequence $\{x_n\}$ by the following iterative scheme:

$$\begin{cases} x_1 \in C \text{ chosen arbitrarily,} \\ C_1 = C, \\ f_\lambda(u_n, y) + \frac{1}{r_n} \langle y - u_n, Ju_n - Jx_n \rangle \geq 0, \quad \forall y \in C, \\ C_{n+1} = \{z \in C_n : \sup_{\lambda \in \Lambda} \phi(z, u_{n,\lambda}) \leq \alpha_n \phi(z, x_1) + (1 - \alpha_n)\phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x \end{cases} \quad (4.4)$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to $\Pi_F x$, where $F = \bigcap_{\lambda \in \Lambda} EP(f_\lambda)$.

Remark 4.8. Throughout the paper, the Mosco convergence plays an important role as the monotone CQ method does in others; see [12,17,18].

References

- [1] E. Blum, W. Oettli, From optimization and variational inequalities to equilibrium problems, *Math. Student* 63 (1994) 123–145.
- [2] S.D. Flam, A.S. Antipin, Equilibrium programming using proximal-link algorithms, *Math. Program.* 78 (1997) 29–41.
- [3] A. Moudafi, M. Thera, Proximal and dynamical approaches to equilibrium problems, in: *Lecture Note in Economics and Mathematical Systems*, vol. 477, Springer-Verlag, New York, 1999, pp. 187–201.
- [4] A. Tada, W. Takahashi, Weak and strong convergence theorems for a nonexpansive mapping and an equilibrium problem, *J. Optim. Theory Appl.* 133 (2007) 359–370.
- [5] Yongfu Su, Meijuan Shang, Xiaolong Qin, An iterative method of solution for equilibrium and optimization problems, *Nonlinear Anal.* 69 (2008) 2709–2719.
- [6] S. Takahashi, W. Takahashi, Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces, *J. Math. Anal. Appl.* 331 (2007) 506–515.
- [7] P.L. Combettes, S.A. Hirstoaga, Equilibrium programming in Hilbert spaces, *J. Nonlinear Convex Anal.* 6 (2005) 117–136.
- [8] L.C. Ceng, J.C. Yao, Hybrid viscosity approximation schemes for equilibrium problems and fixed point problems of infinitely many nonexpansive mappings, *Appl. Math. Comput.* 198 (2008) 729–741.
- [9] L.C. Ceng, J.C. Yao, A hybrid iterative scheme for mixed equilibrium problems and fixed point problems, *J. Comput. Appl. Math.* 214 (2008) 186–201.
- [10] Xiaolong Qin, Yeol Je Cho, Shin Min Kang, Convergence theorems of common elements for equilibrium problems and fixed point problems in Banach spaces, *J. Comput. Appl. Math.* 225 (2009) 20–30.
- [11] P.L. Combettes, S.A. Hirstoaga, Equilibrium programming in Hilbert spaces, *J. Nonlinear Convex Anal.* 6 (2005) 117–136.
- [12] P. Kumam, K. Wattanawitoon, Convergence theorems of a hybrid algorithm for equilibrium problems, *Nonlinear Anal. Hybrid Syst.* (2009) doi:10.1016/j.nahs.2009.02.006.
- [13] K. Takahashi, Zembayashi, Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces, *Nonlinear Anal.* 70 (2009) 45–57.
- [14] W. Takahashi, K. Zembayashi, Strong convergence theorem by a new hybrid method for equilibrium problems and relatively nonexpansive mappings, *Fixed Point Theory Appl.* (2008) doi:10.1155/2008/528476.
- [15] H.K. Xu, An iterative approach to quadratic optimization, *J. Optim. Theory Appl.* 116 (2003) 659–678.
- [16] T. Suzuki, Strong convergence theorems for infinite families of nonexpansive mapping in general Banach spaces, *Fixed Point Theory Appl.* (1) (2005) 103–123.
- [17] Xiaolong Qin, Yeol Je Cho, Shin Min Kang, Haiyun Zhou, Convergence of a modified Halpern-type iteration algorithm for quasi- ϕ -nonexpansive mappings, *Appl. Math. Lett.* 22 (2009) 1051–1055.
- [18] Yongfu Su, Ziming Wang, Hongkun Xu, Strong convergence theorems for a common fixed point of two hemi-relatively nonexpansive mappings, *Nonlinear Anal.* (2009) doi:10.1016/j.na.2009.04.053.

- [19] Yasunori Kimura, Wataru Takahashi, On a hybrid method for a family of relatively nonexpansive mappings in Banach space, *J. Math. Anal. Appl.* 357 (2009) 356–363.
- [20] W. Takahashi, *Convex Analysis and Approximation Fixed points*, Yokohama-Publishers, 2000 (in Japanese).
- [21] S. Matsushita, W. Takahashi, Weak and strong convergence theorems for relatively nonexpansive mappings in Banach spaces, *Fixed Point Theory Appl.* 2004 (2004) 37–47.
- [22] S.Y. Matsushita, W. Takahashi, A strong convergence theorem for relatively nonexpansive mappings in a Banach space, *J. Approx. Theory* 134 (2005) 257–266.
- [23] S. Kamimura, W. Takahashi, Strong convergence of a proximal-type algorithm in a Banach space, *SIAM J. Optim.* 13 (2002) 938–945.
- [24] Yongfu Su, Dongxing Wang, Meijuan Shang, Strong convergence of monotone hybrid algorithm for hemi-relatively nonexpansive mappings, *Fixed Point Theory Appl.* 2008 (2008) 8 pp. Article ID 284613.
- [25] U. Mosco, Convergence of convex sets and of solutions of variational inequalities, *Adv. Math.* 3 (1969) 510–585.
- [26] T. Ibaraki, Y. Kimura, W. Takahashi, Convergence theorems for generalized projections and maximal monotone operators in Banach spaces, *Abstr. Appl. Anal.* (2003) 621–629.
- [27] Shin-ya Matsushita, Wataru Takahashi, A strong convergence theorem for relatively nonexpansive mappings in a Banach space, *J. Approx. Theory* 134 (2005) 257–266.
- [28] Vittorio Colao, Genaro López Acedo, Giuseppe Marino, An implicit method for finding common solutions of variational inequalities and systems of equilibrium problems and fixed points of infinite family of nonexpansive mappings, *Nonlinear Anal.* 71 (2009) 2708–2715.